



Autologous blood transfusion enhances exercise performance - strength of the evidence and physiological mechanisms

Solheim, Sara Amalie; Bejder, Jacob; Breenfeldt Andersen, Andreas; Mørkeberg, Jakob; Nordsborg, Nikolai Baastrup

Published in:
Sports Medicine - Open

DOI:
[10.1186/s40798-019-0204-1](https://doi.org/10.1186/s40798-019-0204-1)

Publication date:
2019

Document version
Publisher's PDF, also known as Version of record

Document license:
[CC BY](#)

Citation for published version (APA):
Solheim, S. A., Bejder, J., Breenfeldt Andersen, A., Mørkeberg, J., & Nordsborg, N. B. (2019). Autologous blood transfusion enhances exercise performance - strength of the evidence and physiological mechanisms. *Sports Medicine - Open*, 5, [30]. <https://doi.org/10.1186/s40798-019-0204-1>

REVIEW ARTICLE

Open Access



Autologous Blood Transfusion Enhances Exercise Performance—Strength of the Evidence and Physiological Mechanisms

Sara Amalie Solheim^{1,2} , Jacob Bejder¹ , Andreas Breenfeldt Andersen¹ , Jakob Mørkeberg² and Nikolai Baastrup Nordsborg^{1*}

Abstract

This review critically evaluates the magnitude of performance enhancement that can be expected from various autologous blood transfusion (ABT) procedures and the underlying physiological mechanisms. The review is based on a systematic search, and it was reported that 4 of 28 studies can be considered of very high quality, i.e. placebo-controlled, double-blind crossover studies. However, both high-quality studies and other studies have generally reported performance-enhancing effects of ABT on exercise intensities ranging from ~70 to 100% of absolute peak oxygen uptake ($\text{VO}_{2\text{peak}}$) with durations of 5–45 min, and the effect was also seen in well-trained athletes. A linear relationship exists between ABT volume and change in $\text{VO}_{2\text{peak}}$. The likely correlation between ABT volume and endurance performance was not evident in the few available studies, but reinfusion of as little as 135 mL packed red blood cells has been shown to increase time trial performance. Red blood cell reinfusion increases endurance performance by elevating arterial oxygen content (CaO_2). The increased CaO_2 is accompanied by reduced lactate concentrations at submaximal intensities as well as increased $\text{VO}_{2\text{peak}}$. Both effects improve endurance performance. Apparently, the magnitude of change in haemoglobin concentration ([Hb]) explains the increase in $\text{VO}_{2\text{peak}}$ associated with ABT because blood volume and maximal cardiac output have remained constant in the majority of ABT studies. Thus, the arterial-venous O_2 difference during exercise must be increased after reinfusion, which is supported by experimental evidence. Additionally, it remains a possibility that ABT can enhance repeated sprint performance, but studies on this topic are lacking. The only available study did not reveal a performance-enhancing effect of reinfusion on 4 × 30 s sprinting. The reviewed studies are of importance for both the physiological understanding of how ABT interacts with exercise capacity and in relation to anti-doping efforts. From an anti-doping perspective, the literature review demonstrates the need for methods to detect even small ABT volumes.

Keywords: Phlebotomy, Transfusion, Arterial oxygen content, Maximal oxygen uptake, Maximal cardiac output, Anti-doping

Key Points

- Autologous blood transfusion increases exercise performance over a broad range of exercise intensities and durations.
- The exact relationship between performance increase and transfusion volume cannot be established due to a lack of studies. Even low-

volume (135 mL) transfusions increase exercise performance.

- Autologous blood transfusion increases maximal oxygen uptake corresponding to the augmented haemoglobin concentration, whereas maximal cardiac output appears unaffected.

Background

One of the first studies of how autologous blood transfusions (ABTs) affect exercise performance was completed in 1960 and demonstrated that sustainable power output at a heart rate of 170 bpm is impaired by ~8% after

* Correspondence: nbn@nexs.ku.dk

¹Department of Nutrition, Exercise and Sport Sciences, University of Copenhagen, Universitetsparken 13, 2100 Copenhagen, Denmark
Full list of author information is available at the end of the article

phlebotomy of ~ 600 mL and increases by ~ 4% after re-infusion of the refrigerated whole blood [1]. Since then, several studies have confirmed ABT to enhance exercise performance in various modes and durations [1–16]. However, ABT is associated with health risks (i.e. bacterial contamination, haemolysis, and allergic reactions) [17] and is against the spirit of sport. Therefore, the use of ABT is prohibited in competitive sports [18] and has been since 1986.

The hitherto-performed ABT studies have never been critically reviewed with a strong focus on research methodology, although the performance-enhancing effect of ABT appears well established and is in agreement with solid physiological principles. Important factors for study interpretation include number of participants, inclusion of controlled trials, blinding, transfusion volume, storage time, storage method, and exercise protocol. Considering the strengths and weaknesses of the existing studies is important not only for the physiological understanding of how ABT interacts with exercise capacity but also for anti-doping efforts, where knowledge of expected effects helps target and prioritize in-competition and out-of-competition testing. Thus, the present review aims to evaluate the magnitude of performance enhancement from various ABT procedures as well as to address the existing knowledge of the associated physiological mechanisms.

A systematic search for peer-reviewed publications was performed at <http://www.pubmed.com> using the following query: (((autologous blood transfusion) OR (red blood cell transfusion)) AND (performance OR exercise)). In addition, the retrieved reference lists were examined for unidentified studies. The literature search was terminated in January 2019. The exclusion criteria were studies published as abstracts only, studies on patients, and studies investigating the combination of ABT and high altitude acclimatization because altitude acclimatization may enhance performance [19]. In total, 28 research articles were included (Table 1).

Effect of ABT on Human Exercise Performance Intensity, Duration, and Training Status

The following section considers the importance of exercise intensity, exercise duration, and participants' training status for the effect of ABT on human performance. In one of the first studies to address ABT's effect on human exercise performance, time to exhaustion (TTE) lasting ~ 5 min increased 23% following reinfusion with 800 mL refrigerated packed red blood cells (RBCs) collected 4 weeks earlier [13]. The TTE was determined from treadmill running at a constant grade with a speed requiring maximal oxygen uptake ($\text{VO}_{2\text{peak}}$). Likewise, TTE lasting ~ 6 min increased by ~ 25% compared to before collection when four participants received cold-

stored packed RBCs from the sequential donation of 1200 mL of blood obtained ~ 30 days earlier [13]. However, the small sample size, the lack of a control group, and the non-blinded design preclude firm conclusions. When applying a double-blind, placebo-controlled crossover design and 7–11 weeks cryopreservation, performance was enhanced by ~ 30% in a TTE lasting ~ 10 min [10] and by 3% in 5 mile and 10 km time trials (TT) [9, 14] in well-trained males receiving a reinfusion of RBCs reconstituted to normal haematocrit (Hct) from a ~ 900 mL donation. Generally, the high-quality ABT studies report enhanced performance by ~ 3–5% compared with pre-reinfusion performance in time trials lasting ~ 30–45 min [7, 9, 14] and that TTE is prolonged by ~ 30% in a test lasting ~ 10 min [10]. It is of note that all the best controlled studies investigated well-trained male athletes (Table 1). Moreover, the performance enhancement was evident when evaluated a few days before and after reinfusion [1, 3–11, 13–16]. Importantly, both reinfusion of cryopreserved RBCs and reinfusion of cold-stored RBCs improve exercise performance irrespective of total haemoglobin mass recovery, which may be incomplete, especially in the context of cold storage, where ≤ 4 weeks elapse between donation and reinfusion [1, 6–8, 13, 30]. In support of the performance-enhancing effect of ABT, both high-quality and low-quality studies generally report that blood reinfusion improves exercise performance in tasks lasting from ~ 5 to ~ 45 min, which corresponds to exercise intensities of approximately 70 to ~ 100% of $\text{VO}_{2\text{peak}}$. Furthermore, performance enhancement has been observed in both untrained and well-trained men (Table 1). Notably, the only study enrolling only women [5] also reported improved performance, but a lack of studies addressing a possible sex-specific effect is apparent. In relative terms, the effect of ABT in open-ended performance tests, i.e. TTE tests with a relatively short duration, appears superior, with an improvement range of 16–40%, to closed-ended tests, i.e. TT tests of longer duration, with an improvement range of 3–5% (Table 1).

ABT Volume and Physiological Effects

In the present section, it is considered whether a dose-dependent effect of ABT exists on maximal aerobic power and exercise performance.

Standard-Volume, High-Volume, and Low-Volume ABT

Reinfusion of cold-stored RBCs from 450–500 mL donations, herein termed 'standard-volume ABT', enhances 3 km treadmill time-trial performance by ~ 5% in moderately trained men [6] and TTE at maximal aerobic workload by ~ 40% in untrained men [16]. However, in some studies, reinfusion of 450 mL whole blood or 200 mL packed RBCs reconstituted with

Table 1: Summary of autologous blood transfusion (ABT) studies examining changes in aerobic power and endurance performance following ABT

Study	Participants (n), training status, and age (years; mean \pm SD)	Study design	ABT volume of whole blood or equivalent when reinfused as packed RBCs (mL storage type, storage duration)	Change in $\dot{V}O_{2peak}$	Performance test (duration/length)	Change in performance
Gullbring et al. [1]	6 M, UT, 34 \pm 8	No control group	530–689 ^{R, 1}	N/A	PWC/TTE cycling (N/A)	4%
Robinson et al. [20]	6 M, UT, 26 \pm 6	No control group	1000–1200 ^{R, 2}	\leftrightarrow	N/A	N/A
Eklom et al. [13]	7 M, WT, 24 \pm 1	No control group	800–1200 ^{R, 4}	\uparrow 9% (1)	TTE running (~ 5 min)	\uparrow 23% (1)
Williams et al. [21]	20 M, MT-WT, 22 \pm 4	Double-blind, placebo-controlled	500 ^{R, 3}	N/A	TTE running (~ 10 min)	\leftrightarrow
Von Rost et al. [8]	6 M, MT, N/A	Controlled	900 ^{R, 3–4}	\uparrow 6%	TTE running (~ 5 min)	\uparrow 37%
Eklom et al. [22]	5 M, WT, 23–31	No control group	800 ^{R, 5}	\uparrow 8%	N/A	N/A
Bell et al. [23]	15 M, MT, 18–24	Controlled	500 ^{R, 3}	\leftrightarrow	TTE cycling (~ 15 min)	\leftrightarrow
Kots et al. [16]	10 M, UT-WT, 22–32	No control group	500 ^{R, 3}	\leftrightarrow	PWC/TTE cycling (~ 10 min)	\uparrow 40%
Williams et al. [24]	16 M, WT, 32 \pm 7	Double-blind, placebo-controlled	460 ^{F, 3}	N/A	TTE running (~ 45 min)	\leftrightarrow
Buick et al. [10]	11 M, WT, 21 \pm 3	Double-blind, placebo-controlled, crossover	900 ^{F, 7}	\uparrow 5%	TTE running (7–10 min)	\uparrow 31–35%
Williams et al. [14]	12 M, WT, 33 \pm 6	Double-blind, placebo-controlled, crossover	920 ^{F, 7}	N/A	TT running (5 miles)	\uparrow 3%
Thomson et al. [25]	4 M, MT-WT, 23 \pm 1	No control group	1000 ^{F, 12}	\uparrow 12%	N/A	N/A
Thomson et al. [26]	4 M, MT-WT, 23 \pm 1	No control group	1000 ^{F, 12}	\uparrow 13%	N/A	N/A
Kanstrup and Eklom [11]	5 M, WT, 25 \pm 4	No control group	900 ^{R, 5}	\uparrow 7%	TTE running (5–6 min)	\uparrow 24%
Robertson et al. [5]	9 W, UT-WT, 23 \pm 2	Single-blind, placebo-controlled, crossover	475 ^{F, 9}	\uparrow 10%	PWC cycling	\uparrow 23%
Celsing et al. [27]	9 M, WT, 29 \pm 2	No control group	3500 ^{R, F, 1–8}	\uparrow 19%	N/A	N/A
Spriet et al. [28]	4 M, WT, 25 \pm 1	No control group	1350 ^{F, 9–11}	\uparrow 7%	N/A	N/A
Muza et al. [29]	12 M, MT-WT, 27 \pm 6	Double-blind, placebo-controlled	600 ^{F, 6}	\uparrow 11%	N/A	N/A
Berglund and Henningsson [2, 30]	11 M, 1 W, WT, 19–35	Single-blind, placebo-controlled	1350 ^{R, 4}	N/A	TT cross-country skiing (15 km)	\uparrow 5%
Brien and Simon [9]	6 M, WT, 24 \pm 5	Double-blind, placebo-controlled, crossover	900 ^{F, 11}	N/A	TT running (10 km)	\uparrow 3%
Celsing et al. [31]	8 M, UT-WT, 28 \pm 3	Controlled	2250 ^{F, 5–7}	\uparrow 7%	N/A	N/A
Sawka et al. [32]	9 M, MT, 30 \pm 7	Double-blind, placebo-controlled	600 ^{F, N/A}	\uparrow 11%	N/A	N/A
Eklom and Berglund [12]	15 M, MT-WT, 27 \pm 3	No control group	1350 ^{F, 8}	\uparrow 8%	TTE running (7–9 min)	\uparrow 16%
Turner et al. [15]	7 M, MT, 26 \pm 3	No control group	900 ^{F, 8–12}	\uparrow 6%	TT cycling (30 min)	\uparrow 5%

Table 1: Summary of autologous blood transfusion (ABT) studies examining changes in aerobic power and endurance performance following ABT (Continued)

Study	Participants (n), training status, and age (years; mean \pm SD)	Study design	ABT volume of whole blood or equivalent when reinfused as packed RBCs (mL storage type, storage duration)	Change in $\text{VO}_{2\text{peak}}$	Performance test (duration/length)	Change in performance
Ziegler et al. [6]	8 M, UT-WT, 33 \pm 2	No control group	450 ^R , 4	\uparrow 5%	TT running (3 km)	\uparrow 5%
Malm et al. [3]	17 M, MT-WT, 34 \pm 8	Placebo-controlled	900 ^F , 15–16	\uparrow 12%	TTE running (6–7 min)	\uparrow 17%
Bennett-Guerrero et al. [4]	4 M, UT-WT, 34 \pm 5 1 week, 29 \pm 7 6 weeks	No control group	900 ^R , 1 or 6	\uparrow 9% 1 week, \uparrow 2% 6 weeks	TTE cycling (12–16 min)	\uparrow 8% 1 week, \downarrow 3% 6 weeks
Bejder et al. [7]	9 M, WT, 29 \pm 5	Double-blind, placebo-controlled, crossover	225 ^R , 4 675 ^R , 4	N/A \leftrightarrow	TT cycling (650 kcal)	\uparrow 4% \uparrow 5%

ABT autologous blood transfusion, M men, W women, $\text{VO}_{2\text{peak}}$ peak oxygen uptake, RBCs red blood cells, R refrigerated, F frozen, \uparrow positive change, \downarrow negative change, \leftrightarrow non-significant change/unchanged, N/A not available, (1) group 1, WT well-trained ($\text{VO}_{2\text{peak}}$ of > 55 mL $\text{O}_2/\text{kg}/\text{min}$), MT moderately trained (45–55 mL $\text{O}_2/\text{kg}/\text{min}$), UT untrained (< 45 mL $\text{O}_2/\text{kg}/\text{min}$), TT time trial, TTE time to exhaustion, PWC physical work capacity given as work (kg-force metres/min) performed to exhaustion

physiological saline to a normal Hct did not result in detectable performance gains in moderately trained to well-trained males, neither following cryopreservation [24, 28] nor after cold storage [21, 23]. Five of eight studies investigating reinfusion of a 'standard-volume' corresponding to ~500 mL reported improved performance, and the three negative findings must be interpreted cautiously (Table 1). There are several possible explanations why some studies have not observed an effect of a single standard-volume autologous reinfusion on $\text{VO}_{2\text{peak}}$ or performance. First, the precision and random variation in the physiological measurements may mask performance-relevant increases. This explanation seems likely since inclusion of, for example, four subjects yields a detection limit for changes in absolute $\text{VO}_{2\text{peak}}$ of ~6% when assuming a measurement error of 3% [33]. Moreover, Bell and co-workers [23] did not observe improved TTE, which appears to be likely related to variation in the small group of 5 subjects. Indeed, a numerical increase of >20% in TTE was apparent in the ABT group, but unfortunately, no data on variation was reported for this outcome variable. Second, RBC storage lesions and degradation after reinfusion can be of importance since only ~70% (cold-stored) or ~50% (cryopreserved) of the haemoglobin (Hb) is preserved after storage [34]. Furthermore, ~25% of the infused RBCs can be degraded within 24 h [35], and a detectable effect of ABT on exercise performance immediately after reinfusion may disappear within a few days. In addition to variations in measures of physiological function and RBC storage lesions, low statistical power for determining subtle performance fluctuations and the selection of time points for performance evaluation are important. For instance, comparison of performance before blood donation and after reinfusion separated by 21 days [21] introduces variation over time as well as a risk of insufficient haemoglobin mass recovery [6, 36]. Indeed, when performance was evaluated prior to phlebotomy and after the reinfusion 21 days later, 10 of 15 participants performed 4–5% worse [21]. However, this finding appears more likely to be related to an insufficient total haemoglobin (tHb) recovery period and natural fluctuations in exercise capacity than lack of effect of ABT on performance. Clearly, variation in measurements, physiological ability, RBC storage lesions, and RBC degradation as well as study design impact the outcome of studies evaluating the effect of standard-volume ABTs on exercise performance.

The same study design factors as outlined above affect studies of ABT with larger volumes, but the expected higher 'signal-to-noise' ratio reduces the risk of not identifying meaningful effects. However, when interpreting

studies of ABT volumes of ~900 mL or more, which is the case for 17 of the studies listed in Table 1, it must be considered whether the associated increase in blood viscosity lowers maximal cardiac output (Q_{max}) more than increased arterial oxygen content ($C_a\text{O}_2$) can compensate for [28] or if it even reduces tissue perfusion. Compromised Q_{max} appears of little relevance because an autosomal dominant erythrocytosis carrier (i.e. increased sensitivity to EPO) with a haematocrit of more than 60% is also an endurance sport Olympic gold medallist [37]. With regard to tissue oxygen delivery, O_2 consumption and contractile power of blood-perfused canine gastrocnemius muscles are compromised only when haematocrit is higher than 60%, despite reduced absolute blood flow [38]. In support of the above observations, exercise performance in rodents is optimal at a haematocrit of approximately 60% [39]. Thus, even markedly elevated blood viscosity after high-volume ABT does not counteract the performance-enhancing effect. This notion is supported by the high-quality studies of ABT volumes ~900 mL reporting improved exercise performance in humans [9, 10, 14].

The possible performance-enhancing effects of ABTs of less than 450 mL, termed 'low-volume ABT' in the present review, have not yet been fully scientifically addressed (Table 1), even though dishonest athletes are suspected to reinfuse low (≤ 150 –200 mL) volumes [40]. To our knowledge, we recently completed the first study on the effect of a low-volume ABT using a randomized, double-blind, placebo-controlled crossover design. We demonstrated that reinfusion with ~135 mL of cold-stored RBCs, corresponding to ~225 mL of whole blood, is sufficient to increase mean power by ~5% and reduce elapsed time by ~4% in a 650 kcal cycling TT [7]. Concomitantly, haemoglobin concentration ([Hb]), Hct, and RBC count increased by 3% compared to their levels 3 days before reinfusion. Hence, even low ABT volumes improve exercise performance.

Is There an ABT Dose-Response Relationship?

Only a few studies have directly addressed whether ABT volume is decisive for the magnitude of effect on absolute $\text{VO}_{2\text{peak}}$ and/or exercise performance. One of the earliest studies to investigate the dose-dependency of ABT on $\text{VO}_{2\text{peak}}$ applied stepwise reinfusions separated by 2–7 days [28]. Each reinfusion consisted of freeze-preserved RBCs, reconstituted with physiological saline to a normal Hct, from 450 mL whole blood harvested 9–11 weeks earlier [28]. The direct effect on endurance performance was not evaluated, but the primary outcome, $\text{VO}_{2\text{peak}}$, surprisingly remained unchanged following the first reinfusion in the four well-trained endurance runners. $\text{VO}_{2\text{peak}}$ did, however, increase non-significantly ($p > 0.05$) by ~200 mL O_2/min

(4%) in all four subjects after the second reinfusion, and an additional significant increase to ~ 340 mL O_2 /min (7%) above control was observed after the third reinfusion [28]. Thus, an undetected dose-dependent response may have been apparent. This interpretation is supported by the gradual increase in VO_{2peak} observed with four successive reinfusions of 450 mL [31]. Thus, the impact of ABT on VO_{2peak} apparently follows a dose-response relationship, as could be expected, although a well-designed study to address this question is lacking.

Does Storage Technique Impact ABT's Effect on Exercise Performance?

Two main techniques are used for RBC storage: refrigeration (cold storage) or freezing (cryopreservation). In the clinical setting, cold-stored RBCs must be reinfused after 35–42 days, depending on the storage solution, due to increasing storage lesions, whereas cryopreservation allows for storage for up to 30 years [41]. Cold-stored blood appears to be the least probable strategy used by dopers because 20–59 days is required for full tHb restoration after an ~ 550 mL phlebotomy [42] and the effect on [Hb], and thus performance, may be counteracted by suppressed haemoglobin mass at the time of reinfusion. On the other hand, pre-reinfusion handling of cryopreserved blood is associated with a 20% larger loss of the initially removed Hb compared with that of refrigerated blood due to the necessary freeze-thaw-wash process following cryopreservation [34]. Moreover, cryopreservation is more expensive and technically challenging than cold storage. No studies have directly compared the impact of storage technique on performance. However, it is important to note that the evaluation of ABT's effect on exercise capacity requires that the performance evaluation be completed a few days before and after reinfusion, independent of storage technique. Evidently, both storage techniques are able to enhance aerobic performance (Table 1), although the shortcomings of the existing protocols need to be acknowledged.

Study Design Considerations

Several factors influence the interpretation of ABT and exercise performance studies. These factors include storage technique, number of participants, inclusion of control trials, blinding procedure, transfusion volume, storage time, exercise protocol, and statistical analyses. From a human performance and anti-doping perspective, the lack of a statistically significant difference following a treatment is of limited value if just one participant gains a competitive advantage. The margin of difference between competitors is small, and a $<1\%$ effect is of relevance. However, such small changes are difficult to detect in a scientific study. A useful approach

in this context could be the evaluation of clinical significance or relevance by reporting effect sizes and minimal clinically important differences [43]. However, the sensitivity issue appears to be of less importance in ABT studies, as a performance-enhancing effect of several percent is present, even when low-volume transfusions are applied [7]. To date, six studies with endurance performance measurements have applied a double-blind, placebo-controlled design with a total of 6–20 participants [7, 9, 10, 14, 21, 24]. All except two [21, 24] of these studies reported improved performance after the reinfusion of autologous blood, which may be ascribed to a sample size of 5 [21] or 8 [24] in each group or variation in the measurements.

Physiological Mechanisms Responsible for the Performance-Enhancing Effect of ABT

Theoretically, endurance performance is related to VO_{2peak} relative to body weight and the % of VO_{2peak} that can be sustained during prolonged intense endurance exercise (i.e. 30–40 min for a 10-km run). However, how does ABT alter VO_{2peak} and the ability to sustain a high percentage of VO_{2peak} ? In the following sections, the physiological mechanisms that may explain the performance-enhancing effect of ABT are considered.

ABT Increases Maximal Muscular Oxygen Delivery

Maximal O_2 delivery to the contracting limb muscles can be augmented via increased limb blood flow, increased C_aO_2 , or both. An increase in maximal limb blood flow requires increased Q_{max} or altered blood flow distribution. In the 'Is Maximal Cardiac Output Increased by ABT?' and 'Is Maximal Limb O_2 Delivery and Extraction Increased by ABT?' sections, ABT's potential effect on Q_{max} , limb blood flow, and C_aO_2 is addressed.

Is Maximal Cardiac Output Increased by ABT?

When maximal exercise with a large muscle mass is performed in normoxia, Q_{max} determines the upper limit for O_2 delivery to the active muscle groups and thereby becomes the limiting factor for VO_{2peak} in endurance-trained, untrained, and generally healthy individuals [44]. Hence, increasing Q_{max} will increase an individual's maximal oxygen uptake and thereby also maximal endurance performance by allowing the individual to exercise at a higher absolute workload, while metabolism remains primarily aerobic [45]. In cross-sectional studies, total blood volume and tHb are tightly associated with VO_{2peak} , whereas [Hb] is not [46]. In agreement with this idea, Q_{max} is closely related to VO_{2peak} [47], and thus, it may be hypothesized that an increased blood volume can increase Q_{max} and thereby performance. Indeed, a training-induced expansion of total blood volume appears to be the primary cause of increased

$\text{VO}_{2\text{peak}}$ via increased Q_{max} [48]. Thus, it is tempting to suggest that the ABT-induced increase in tHb increases $\text{VO}_{2\text{peak}}$ by increasing Q_{max} . However, it is important to note that the close association between absolute $\text{VO}_{2\text{peak}}$ and tHb when determined in a mixed group of humans is caused primarily by differences in size, including ventricular mass and total blood volume, which defines Q_{max} [47], whereas no close association is evident in a homogenous group of athletes [49]. With respect to the effect of blood reinfusion on $\text{VO}_{2\text{peak}}$, it is important that total blood volume alterations appear to depend on the time of investigation after reinfusion. Total blood volume appears unaffected ≥ 24 h post-reinfusion of 800 to 2250 mL of refrigerated whole blood [22] as well as refrigerated [50] or freeze-preserved [31, 32, 50] RBCs reconstituted with physiological saline to a normal Hct. In contrast, 1 h after reinfusion of packed RBCs from 900 mL of whole blood, a 5% increase in total blood volume occurs together with increases in Hct (2%), [Hb] (4%), $\text{VO}_{2\text{peak}}$ (7%), and cycling performance lasting 3–6 min (24%) [11]. As such, total blood volume may be unaffected by ABT when determined more than 24 h after reinfusion due to a rapid compensatory reduction in plasma volume. However, the small number of studies on ABT and total blood volume and the variances and challenges associated with blood volume determination preclude a firm conclusion.

Reinfusion of autologous blood acutely increases central venous pressure [20], which usually increases Q_{max} due to an enlarged end-diastolic volume and thereby causes a higher stroke volume and Q_{max} , according to the Frank-Starling mechanism [51]. However, it is likely that the increased central venous pressure after blood reinfusion is transient (< 24 h) and related to the acute volume expansion, but this remains to be investigated. The most direct approach to determine the effect of ABT on Q_{max} is to measure Q_{max} . A few studies have directly evaluated Q_{max} after blood reinfusion. In 7 males who received RBCs obtained from 900 mL donations 2–3 months earlier, no difference existed approximately 24 h after reinfusion in the invasively determined Q_{max} of ~ 22 L/min, despite a clear $\sim 3\%$ increase in $\text{VO}_{2\text{peak}}$ [15]. In 9 women, reinfusion of RBCs from two 450 mL phlebotomies, conducted 9 and 17 weeks prior, did not alter the Q_{max} of ~ 17 L/min, as indirectly determined 48 h later by CO_2 rebreathing [5]. Moreover, in four male endurance runners, the reinfusion of RBCs from 900–1350 mL blood 3–4 weeks after phlebotomy increased $\text{VO}_{2\text{peak}}$ but only non-significantly increased dye dilution-determined Q_{max} from ~ 28 to ~ 33 L/min approximately 2 days later [28]. Thus, Q_{max} appears unaffected even by reinfusion of rather large volumes and cannot explain the increase in $\text{VO}_{2\text{peak}}$ and performance after ABT. In this context, it must be noted that

determination of Q_{max} is technically difficult and associated with variance; therefore, a moderate undetected effect of ABT on Q_{max} may exist.

Is Maximal Limb O_2 Delivery and Extraction Increased by ABT?

Limb blood flow can be as high as 2000–3000 mL/min per kg of muscle [52], and the muscular oxygen uptake capacity is ~ 600 mL/kg/min in healthy subjects, which is much higher than the amount of O_2 supplied during whole-body exercise [53]. Hence, it is evident that limb blood flow is limited by Q_{max} during intense whole-body exercise. Thus, if maximal limb blood flow during intense whole-body exercise is increased by reinfusion, this would increase maximal aerobic energy production and thereby performance. However, maximal limb blood flow appears unlikely to increase since Q_{max} appears unaffected by reinfusion, as highlighted in the 'Is Maximal Cardiac Output Increased by ABT?' section. In this context, it must be noted that, for example, brain blood flow is primarily dependent on C_aO_2 [54], and therefore, the ABT-induced increase in arterial oxygen content may reduce blood flow to other tissues and organs. As a consequence, it remains a possibility that more blood is redistributed to the exercising muscles and that maximal limb blood flow is marginally increased by RBC reinfusion despite unaltered Q_{max} . However, this hypothesis remains to be investigated.

Reinfusion of autologous blood is usually [7, 10, 13, 14, 28, 31, 32], but not always [3], accompanied by elevated Hct and/or [Hb] compared to the pre-reinfusion levels. Accordingly, reinfusion increases the O_2 carrying capacity per litre of blood, and this has been highlighted in previous reviews as the main physiological mechanism causing enhanced performance following ABT [55]. Indeed, stepwise reinfusions of four blood bags, in eight untrained to moderately trained men, resulted in $\text{VO}_{2\text{peak}}$ increases of approximately 40 mL O_2 /min (1%), 130 mL O_2 /min (3%), 230 mL O_2 /min (5%), and 320 mL O_2 /min (7%) and increases in [Hb] of approximately 4%, 7%, 12%, and 13% from the first to the last reinfusion, respectively [31], as mentioned in the 'Is There an ABT Dose-Response Relationship' section. This result resembles the findings of Spriet and co-workers [28] and corresponds to an approximately 0.47% increase in $\text{VO}_{2\text{peak}}$ per 1% increase in [Hb] ($R^2 = 0.97$) [31]. For the ABT-induced increase in [Hb] to augment $\text{VO}_{2\text{peak}}$ without affecting Q_{max} , the arterial-venous O_2 difference (a-v O_2 difference) across the exercising muscle groups must be increased. The ABT-induced increase in C_aO_2 and concurrent unchanged [15, 25] or reduced [22] mixed venous O_2 content (C_vO_2) strongly suggest an increased extraction [15] after reinfusion, but the small sample sizes ($n = 4$ –7) preclude a firm conclusion.

ABT, Anaerobic Energy Production, and Metabolic Acidosis Blood Lactate Response to Exercise

Increased maximal capacity for oxygen delivery likely reduces the anaerobic energy contribution at intensities above $\sim 70\%$ of $\text{VO}_{2\text{peak}}$, as illustrated by reduced plasma lactate concentrations ($[\text{La}^-]$) after ABT measured at the same absolute workload [5]. The reinfusion of RBCs packed saline obtained from phlebotomy of 900 mL and 1350 mL whole blood causes a marked reduction in blood lactate values following exercise at 91% of $\text{VO}_{2\text{peak}}$ (from ~ 6 mM before reinfusion to ~ 2 mM after) and after maximal treadmill running at 100% of control $\text{VO}_{2\text{peak}}$ (from ~ 12 mM before to ~ 5 mM after), concomitant with less disturbed arterial acid-base status 5 min after exercise [28]. Furthermore, blood reinfusion reduces the lactate concentration by approximately 50% in runners when determined 4 min after submaximal ($\sim 70\%$ $\text{VO}_{2\text{peak}}$) treadmill running [10]. In five trained subjects, venesection increased the lactate response to exercise at $\sim 70\%$ of $\text{VO}_{2\text{peak}}$ from 6.0 to 7.6 mM, while reinfusion reduced lactate to 4.6 mM [22]. This reduction in plasma-borne markers of anaerobic energy contribution during submaximal exercise may increase the $\%\text{VO}_{2\text{peak}}$ defining the lactate threshold, which is an important determinant of endurance performance [56, 57]. Since most endurance competitions are not performed at intensities corresponding to 100% of $\text{VO}_{2\text{peak}}$, this may be the most important performance-enhancing effect of ABT. However, it should also be acknowledged that some parts, e.g. the final part of a 5000-m or 10,000-m race, a cycling stage with mountain climbs or the like, are likely to elicit $\text{VO}_{2\text{peak}}$.

Blood pH Response to Exercise

During intense exercise, anaerobic energy production results in increased plasma hydrogen concentration ($[\text{H}^+]$) [58]. Metabolic acidosis is immediately counteracted by hyperventilation, bicarbonate buffering, and non-bicarbonate buffering. The relationship between changes in plasma lactate concentration and pH is indicated by the in vivo buffer capacity: $\Delta[\text{La}^-]/\Delta\text{pH}$ [59]. RBCs are primary extracellular buffers, and therefore, the total blood buffer capacity depends on the total RBC volume and the expression of RBC membrane-bound transport systems, such as the AE1 isoform of the anion-exchange transporter [60, 61] and the MCT1 isoform of the monocarboxylate transporter (La^-/H^+ co-transporter) [61], as well as the enzyme cytoplasmic carbonic anhydrase, which catalyses the reversible hydration of carbon dioxide (CO_2); $\text{CO}_2 + \text{H}_2\text{O}$ (water) \rightleftharpoons H_2CO_3 (carbonic acid) \rightleftharpoons HCO_3^- (bicarbonate) + H^+ [60, 61]. Hence, the RBCs added through ABT have the potential to theoretically improve systemic pH regulation. However, the possible effect is small since a low-volume ABT expands the total RBC volume by only a few percent. Nevertheless, the relevance for performance in

elite sports with small margins may exist, but this remains to be investigated.

Possible Influence of ABT on Anaerobic Performance and Repeated Sprints?

The reinfusion of autologous blood increases the maximal muscular oxygen uptake, which is the product of limb blood flow and arterial-venous O_2 difference. Blood reinfusions might also increase performance in short, anaerobic, intermittent exercises. Since the aerobic energy provision increases when sprints are repeated [62, 63] and the aerobic capacity is important for recovery between sprints [64], it appears likely that repeated sprint ability is improved by blood reinfusion. $[\text{La}^-]$ in blood and muscle increases progressively during a 4×30 s all-out air-braked cycle test in moderately trained males [65]. Additionally, muscle La^- accumulates non-linearly during a 2×30 s all-out cycle test at a work load corresponding to $0.075 \times \text{kg}$ of body weight [66] and during 3×30 s all-out isokinetic cycling at 100 rpm in healthy, recreationally active subjects [58]. The highest impact of blood reinfusion on repeated sprint ability is thus likely to occur in the later sprints. In support of this notion, the level of aerobic ATP provision during sprints has been found to increase progressively as work duration increases [67, 68] and as sprints are repeated [62, 63]. Additionally, the potentially improved blood buffer capacity during polycythaemia might contribute to the extracellular pH defence against lactic acidosis, as outlined in the 'ABT, Anaerobic Energy Production, and Metabolic Acidosis' section [59, 61]. Furthermore, the nature of the recovery between sprints is likely of importance for repeated sprint ability. For example, active recovery is advantageous for the mean power output of sprints when repeated Wingate anaerobic tests are performed [62]. The O_2 availability for mitochondria is known to be decisive for creatine phosphate recovery [69], and an ABT-induced increase in C_aO_2 thus appears likely to improve recovery between sprints. Supporting this theory, the 11% increase in $[\text{Hb}]$, following recombinant human EPO administration, reduces the accumulation of blood La^- and plasma hypoxanthine after fifteen 6-s high-intensity uphill running sprints with 24 s of passive rest [70]. Interestingly, in the only study to investigate the effect of ABT on repeated cycling sprint ability, we observed no change in performance after reinfusion, despite an $\sim 9\%$ increase in tHb and $[\text{Hb}]$ [7]. The combined effect of the reinfusion of ~ 135 and ~ 235 mL cold-stored RBCs, corresponding to approximately 675 mL whole blood, did not alter 4×30 s all-out cycling performance, either when determined as the mean power of all sprints or when evaluated as the mean power in each sprint. The two reinfusions were performed within ~ 24 h, and the performance was evaluated a few days before the first reinfusion and after the last

reinfusion, separated by 8 days, as opposed to before the phlebotomy and after the reinfusion. This allowed for the evaluation of ABT-induced changes, although the haematological parameters had not returned to the pre-phlebotomy level at the time of reinfusion. The results suggest that reinfusion with ~370 mL of packed RBCs does not affect repeated sprint ability as measured by 4 × 30 s of all-out cycling interspersed with 4 min of rest. However, the effect of ABT in other repeated sprint modalities and the effect of larger ABT volumes are unknown.

Overall, the literature is scarce regarding the effects of ABT on repeated sprint performance. Nevertheless, the aforementioned potential effects of ABT in enhancing O₂ delivery and blood buffer capacity might be reflected in improvement in fatigue resistance during repeated sprints and thereby enhanced repeated sprint performance in some repeated sprint modalities.

Conclusion

The literature review revealed that ABT enhances performance across a span of exercise intensities from ~70 to 100% of VO_{2peak} and durations of 5–45 min, and these results were also observed in well-trained athletes. The impact of ABT on VO_{2peak} apparently exhibits a dose-response relationship, but well-designed studies of likely associated gradual performance effects are lacking. However, reinfusion with as little as 135 mL packed RBCs increases time-trial performance. Red blood cell reinfusion increases endurance performance by elevating C_aO₂. The increased C_aO₂ is accompanied by increased VO_{2peak} as well as reduced lactate concentrations, i.e. reduced anaerobic energy contribution at submaximal intensities. Both effects improve endurance performance, the latter because most endurance competitions are not performed at intensities corresponding to 100% of VO_{2peak}. Apparently, the magnitude of change in haemoglobin concentration explains the increase in VO_{2peak} associated with ABT because blood volume and maximal cardiac output remain constant in the majority of ABT studies. Thus, the arterial-venous O₂ difference during exercise must be increased after reinfusion, which is supported by experimental evidence. Additionally, it remains a possibility that ABT can enhance repeated sprint performance, but studies are lacking. The only available study did not observe a performance-enhancing effect of reinfusion on 4 × 30 s sprinting.

The findings are of importance for both the physiological understanding of how ABT interacts with exercise capacity and in relation to anti-doping efforts. From an anti-doping perspective, the literature review demonstrates the need for analytical methods able to detect even minor blood manipulations.

Abbreviations

[Hb]: Haemoglobin concentration; [La⁻]: Lactic acid concentration; ABT: Autologous blood transfusion; a-v O₂ difference: Arterial-venous O₂ difference; C_aO₂: Arterial oxygen content; CO₂: Carbon dioxide; C_vO₂: Mixed venous oxygen content; EPO: Erythropoietin; H⁺: Hydrogen ion; H₂CO₃: Carbonic acid; H₂O: Water; Hb: Haemoglobin; HCO₃⁻: Bicarbonate; Hct: Haematocrit; La⁻: Lactic acid; Q_{max}: Maximal cardiac output; RBC: Red blood cell; tHb: Total haemoglobin mass; TT: Time trial; TTE: Time to exhaustion; VO_{2peak}: Peak oxygen uptake

Acknowledgements

Not applicable.

Authors' Contributions

SAS and NBN planned the review. SAS retrieved the relevant literature. All authors engaged in drafting and revising the manuscript. All authors approved the final version of the manuscript.

Funding

During the preparation of this review, Sara Amalie Solheim was funded by Anti Doping Denmark and the Partnership for Clean Competition, Jacob Bejder by the Partnership for Clean Competition, Andreas Breenfeldt Andersen by World Anti-Doping Agency, and Jakob Mørkeberg served as Senior Science Manager at Anti Doping Denmark.

Availability of Data and Materials

Not applicable. The data that support the findings of this review are available from the cited research articles.

Ethics Approval and Consent to Participate

Not applicable.

Consent for Publication

Not applicable.

Competing Interests

The authors, Sara Amalie Solheim, Jacob Bejder, Andreas Breenfeldt, Jakob Mørkeberg, and Nikolai Baastrup Nordsborg, declare that they have no competing interests.

Author details

¹Department of Nutrition, Exercise and Sport Sciences, University of Copenhagen, Universitetsparken 13, 2100 Copenhagen, Denmark. ²Anti Doping Denmark, Brøndby, Denmark.

Received: 20 November 2018 Accepted: 25 June 2019

Published online: 08 July 2019

References

1. Gullbring B, Holmgren A, Sjostrand T, Strandell T. The effect of blood volume variations on the pulse rate in supine and upright positions and during exercise. *Acta Physiol Scand*. 1960;50:62–71.
2. Berglund B, Hemmingsson P. Effect of reinfusion of autologous blood on exercise performance in cross-country skiers. *Int J Sports Med*. 1987;8:231–3.
3. Malm CB, Khoo NS, Granlund I, Lindstedt E, Hult A. Autologous doping with cryopreserved red blood cells - effects on physical performance and detection by multivariate statistics. *PLoS One*. 2016;11:e0156157.
4. Bennett-Guerrero E, Lockhart EL, Bandarenko N, Campbell ML, Natoli MJ, Jamnik VK, et al. A randomized controlled pilot study of VO₂ max testing: a potential model for measuring relative in vivo efficacy of different red blood cell products. *Transfusion*. 2017;57:630–6.
5. Robertson RJ, Gilcher R, Metz KF, Caspersen CJ, Allison TG, Abbott RA, et al. Hemoglobin concentration and aerobic work capacity in women following induced erythrocythemia. *J Appl Physiol*. 1984;57:568–75.
6. Ziegler AK, Grand J, Stangerup I, Nielsen HJ, Dela F, Magnussen K, et al. Time course for the recovery of physical performance, blood hemoglobin, and ferritin content after blood donation. *Transfusion*. 2015;55:898–905.
7. Bejder J, Andersen AB, Solheim SA, Gybel-Brask M, Secher NH, Johansson PI, et al. Time trial performance is sensitive to low-volume autologous blood transfusion. [Epub ahead of print]. *Med Sci Sports Exerc*. 2018. <https://doi.org/10.1249/MSS.0000000000001837>.

8. Von Rost R, Hollmann W, Liesen H, Schultzen D. Über den einfluss einer erythrozyten-retransfusion auf die kardio-pulmonale leistungsfähigkeit. *Sportarzt u. Sportmed.* 1975;26:137–44.
9. Brien AJ, Simon TL. The effects of red blood cell infusion on 10-km race time. *JAMA.* 1987;257:2761–5.
10. Buick FJ, Gledhill N, Froese AB, Spriet L, Meyers EC. Effect of induced erythrocythemia on aerobic work capacity. *J Appl Physiol.* 1980;48:636–42.
11. Kanstrup IL, Ekblom B. Blood volume and hemoglobin concentration as determinants of maximal aerobic power. *Med Sci Sports Exerc.* 1984;16:256–62.
12. Ekblom BT, Berglund B. Effect of erythropoietin administration on maximal aerobic power. *Scand J Med Sci Sport.* 1991;1:88–93.
13. Ekblom BT, Goldbarg AN, Gullbring B. Response to exercise after blood loss and reinfusion. *J Appl Physiol.* 1972;33:175–80.
14. Williams MH, Wesseldine S, Somma T, Schuster R. The effect of induced erythrocythemia upon 5-mile treadmill run time. *Med Sci Sports Exerc.* 1981;13:169–75.
15. Turner DL, Hoppeler H, Noti C, Gurtner HP, Gerber H, Schena F, et al. Limitations to VO₂max in humans after blood retransfusion. *Respir Physiol.* 1993;92:329–41.
16. Kots YM, Shcherba MM, Kolker YS, Gorodetskii VD, Sin LD. Experimental study of the relationship between the blood hemoglobin concentration and physical aerobic working capacity. *Hum Physiol.* 1976;4:43–9.
17. Domen RE. Adverse reactions associated with autologous blood transfusion: evaluation and incidence at a large academic hospital. *Transfusion.* 1998;38:296–300.
18. World Anti-Doping Agency. The 2018 Prohibited List. Montreal, Quebec: World Anti-Doping Agency; 2017.
19. Chapman RF, Laymon AS, Levine BD. Timing of arrival and pre-acclimatization strategies for the endurance athlete competing at moderate to high altitudes. *High Alt Med Biol.* 2013;14:319–24.
20. Robinson BF, Epstein SE, Kahler RL. Circulatory effects of acute expansion of blood volume: studies during maximal exercise and at rest. *Circ Res.* 1966;XIX:26–33.
21. Williams MH, Goodwin AR, Perkins R, Bocrie J. Effect of blood reinjection upon endurance capacity and heart rate. *Med Sci Sports Exerc.* 1973;5:181–6.
22. Ekblom BT, Wilson G, Åstrand PO. Central circulation during exercise after venesection and reinfusion of red blood cells. *J Appl Physiol.* 1976;40:379–83.
23. Bell RD, Card RT, Johnson MA, Cunningham TA, Baker F. Blood doping and athletic performance. *Aust J Sport Med.* 1976;8:133–9.
24. Williams MH, Lindhjem M, Schuster R. The effect of blood infusion upon endurance capacity and ratings of perceived exertion. *Med Sci Sports Exerc.* 1978;10:113–8.
25. Thomson JM, Stone J, Ginsburg A, Hamilton P. O₂ transport during exercise following blood reinfusion. *J Appl Physiol Respir Env Exerc Physiol.* 1982;53:1213–9.
26. Thomson JM, Stone JA, Ginsburg AD, Hamilton P. The effects of blood reinfusion during prolonged, heavy exercise. *Can J Appl Sport Sci.* 1983;8:72–8.
27. Celsing F, Nystrom J, Pihlstedt P, Werner B, Ekblom B. Effect of long-term anemia and retransfusion on central circulation during exercise. *J Appl Physiol.* 1986;61:1358–62.
28. Spriet LL, Gledhill N, Froese AB, Wilkes DL. Effect of graded erythrocythemia on cardiovascular and metabolic responses to exercise. *J Appl Physiol.* 1986;61:1942–8.
29. Muza SR, Sawka MN, Young AJ, Dennis RC, Gonzalez RR, Martin JW, et al. Elite special forces: physiological description and ergogenic influence of blood reinfusion. *Aviat Space Environ Med.* 1987;58:1001–4.
30. Berglund B, Hemmingsson P, Birgegård G. Detection of autologous blood transfusion in cross-country skiers. *Int J Sports Med.* 1987;8:66–70.
31. Celsing F, Svedenahag J, Pihlstedt P, Ekblom B. Effects of anaemia and stepwise-induced polycythemia on maximal aerobic power in individuals with high and low haemoglobin concentrations. *Acta Physiol Scand.* 1987;129:47–54.
32. Sawka MN, Dennis RC, Gonzalez RR, Young AJ, Muza SR, Martin JW, et al. Influence of polycythemia on blood volume and thermoregulation during exercise-heat stress. *J Appl Physiol.* 1987;62:912–8.
33. Pettitt RW, Clark IE, Ebner SM, Sedgeman DT, Murray SR. Gas exchange threshold and VO₂max testing for athletes. *J Strength Cond Res.* 2013;27:549–55.
34. Ashenden M, Mørkeberg J. Net haemoglobin increase from reinfusion of refrigerated vs. frozen red blood cells after autologous blood transfusions. *Vox Sang.* 2011;101:320–6.
35. Luten M, Roerdinkholder-Stoelwinder B, Schaap NPM, De Grip WJ, Bos HJ, Bosman GJCGM. Survival of red blood cells after transfusion: a comparison between red cells concentrates of different storage periods. *Transfusion.* 2008;48:1478–85.
36. Pottgiesser T, Specker W, Umhau M, Dickhuth H-H, Roecker K, Schumacher YO. Recovery of hemoglobin mass after blood donation. *Transfusion.* 2008;48:1390–7.
37. Juvonen E, Ikkala E, Fyhrquist F, Ruutu T. Autosomal dominant erythrocytosis caused by increased sensitivity to erythropoietin. *Blood.* 1991;78:3066–9.
38. Gaehetgens P, Kreutz F, Albrecht KH. Optimal hematocrit for canine skeletal muscle during rhythmic isotonic exercise. *Eur J Appl Physiol Occup Physiol.* 1979;41:27–39.
39. Schuler B, Arras M, Keller S, Rettich A, Lundby C, Vogel J, et al. Optimal hematocrit for maximal exercise performance in acute and chronic erythropoietin-treated mice. *Proc Natl Acad Sci U S A.* 2010;107:419–23.
40. Marty D, Nicholson P, Haas U. Report to the President of the Union Cycliste Internationale. Lausanne: Union Cycliste Internationale; 2015.
41. European Directorate for the Quality of Medicines & HealthCare of the Council of Europe. Component monographs. Part B. Red cell components. 19th edition Guid to Prep use Qual Assur blood components. Strasbourg; 2017. p. 289–323.
42. Pottgiesser T, Specker W, Umhau M, Dickhuth HH, Roecker K, Schumacher YO. Recovery of hemoglobin mass after blood donation. *Transfusion.* 2008;48:1390–7.
43. Page P. Beyond statistical significance: clinical interpretation of rehabilitation research literature. *Int J Sports Phys Ther.* 2014;9:726–36.
44. Saltin B, Strange S. Maximal oxygen uptake: “old” and “new” arguments for a cardiovascular limitation. *Med Sci Sports Exerc.* 1992;24:30–7.
45. Joyner MJ, Coyle EF. Endurance exercise performance: the physiology of champions. *J Physiol.* 2008;586:35–44.
46. Montero D, Lundby C. Regulation of red blood cell volume with exercise training. *Compr Physiol.* 2018;9:149–64.
47. Lundby C, Montero D, Joyner M. Biology of VO₂ max: looking under the physiology lamp. *Acta Physiol (Oxf).* 2017;220:218–28.
48. Bonne TC, Doucende G, Flück D, Jacobs RA, Nordsborg NB, Robach P, et al. Phlebotomy eliminates the maximal cardiac output response to six weeks of exercise training. *Am J Physiol Integr Comp Physiol.* 2014;306:R752–60.
49. Bejder J, Bonne TC, Nyberg M, Sjøberg KA, Nordsborg NB. Physiological determinants of elite mountain bike cross-country Olympic performance. [Epub ahead of print]. *J Sports Sci.* 2018. <https://doi.org/10.1080/02640414.2018.1546546>.
50. Mørkeberg J, Belhage B, Rasmussen AB, Damsgaard R, Ashenden MJ, Dziegiel MH. Changes in red blood cell volume, plasma volume, and total blood volume after autologous blood collections. *Transfusion.* 2008;48:1524–6.
51. Levine BD, Lane LD, Buckley JC, Friedman DB, Blomqvist CG. Left ventricular pressure-volume and Frank-Starling relations in endurance athletes. Implications for orthostatic tolerance and exercise performance. *Circulation.* 1991;84:1016–23.
52. Saltin B, Radegran G, Koskolou MD, Roach RC. Skeletal muscle blood flow in humans and its regulation during exercise. *Acta Physiol Scand.* 1998;162:421–36.
53. Poole DC, Richardson RS. Determinants of oxygen uptake. Implications for exercise testing. *Sports Med.* 1997;24:308–20.
54. Mühling J, Dehne MG, Sablotzki A, Hempelmann G. Cerebral blood flow velocity during isovolemic hemodilution and subsequent autologous blood retransfusion. *Can J Anesth.* 1999;46:550–7.
55. Jones M, Tunstall Pedoe DS. Blood doping - a literature review. *Br J Sports Med.* 1989;23:84–8.
56. Bassett DR, Howley ET. Limiting factors for maximum oxygen uptake and determinants of endurance performance. *Med Sci Sports Exerc.* 2000;32:70–84.
57. Coyle EF, Coggan AR, Hopper MK, Walters TJ. Determinants of endurance in well-trained cyclists. *J Appl Physiol.* 1988;64:2622–30.
58. Spriet LL, Lindinger MI, McKelvie RS, Heigenhauser GJ, Jones NL. Muscle glycogenolysis and H⁺ concentration during maximal intermittent cycling. *J Appl Physiol.* 1989;66:8–13.

59. Böning D, Maassen N, Thomas A, Steinacker JM. Extracellular pH defense against lactic acid in normoxia and hypoxia before and after a Himalayan expedition. *Eur J Appl Physiol*. 2001;84:78–86.
60. Swietach P, Tiffert T, Mauritz JMA, Seear R, Esposito A, Kaminski CF, et al. Hydrogen ion dynamics in human red blood cells. *J Physiol*. 2010;588:4995–5014.
61. Juel C, Lundby C, Sander M, Calbet JAL, van Hall G. Human skeletal muscle and erythrocyte proteins involved in acid-base homeostasis: adaptations to chronic hypoxia. *J Physiol*. 2003;548:639–48.
62. Lopez E-ID, Smoliga JM, Zavorsky GS. The effect of passive versus active recovery on power output over six repeated wingate sprints. *Res Q Exerc Sport*. 2014;85:519–26.
63. McGawley K, Bishop DJ. Oxygen uptake during repeated-sprint exercise. *J Sci Med Sport*. 2015;18:214–8.
64. Bogdanis GC, Nevill ME, Boobis LH, Lakomy HK, Nevill AM. Recovery of power output and muscle metabolites following 30 s of maximal sprint cycling in man. *J Physiol*. 1995;482(Pt 2):467–80.
65. Hargreaves M, McKenna MJ, Jenkins DG, Warmington SA, Li JL, Snow RJ, et al. Muscle metabolites and performance during high-intensity, intermittent exercise. *J Appl Physiol*. 1998;84:1687–91.
66. Bogdanis GC, Nevill ME, Boobis LH, Lakomy HK. Contribution of phosphocreatine and aerobic metabolism to energy supply during repeated sprint exercise. *J Appl Physiol*. 1996;80:876–84.
67. Calbet JAL, Chavarren J, Dorado C. Fractional use of anaerobic capacity during a 30- and a 45-s Wingate test. *Eur J Appl Physiol Occup Physiol*. 1997;76:308–13.
68. Medbø JJ, Tabata I. Relative importance of aerobic and anaerobic energy release during short-lasting exhausting bicycle exercise. *J Appl Physiol*. 1989;67:1881–6.
69. Hogan MC, Richardson RS, Haseler LJ. Human muscle performance and PCr hydrolysis with varied inspired oxygen fractions: a 31P-MRS study. *J Appl Physiol*. 1999;86:1367–73.
70. Balsom PD, Ekblom B, Sjodin B. Enhanced oxygen availability during high intensity intermittent exercise decreases anaerobic metabolite concentrations in blood. *Acta Physiol Scand*. 1994;150:455–6.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)